

# On compatibility of Bohmian mechanics with standard quantum mechanics

Hrvoje Nikolić

Theoretical Physics Division, Rudjer Bošković Institute,  
P.O.B. 180, HR-10002 Zagreb, Croatia

E-mail: hrvoje@thphys.irb.hr

February 1, 2008

## Abstract

It is shown that the apparent incompatibility of Bohmian mechanics with standard quantum mechanics, found by Akhavan and Golshani [quant-ph/0305020], is an artefact of the fact that the initial wavefunction they use, being proportional to a  $\delta$ -function, is not a regular wavefunction.

PACS numbers: 03.65.Ta

In [1], Golshani and Akhavan argued that Bohmian mechanics [2] was not compatible with standard quantum mechanics. They studied a case in which a 2-particle wavefunction  $\psi(x_1, y_1, x_2, y_2, t)$  describing a pair of entangled particles moving in the  $x$ - $y$  plane was such that the Bohmian trajectories of the particles obey

$$y_1(t) + y_2(t) = 0, \quad (1)$$

provided that the initial positions of the particles obey

$$y_1(0) + y_2(0) = 0. \quad (2)$$

They have shown that (1) is incompatible with standard quantum mechanics and suggested that this could be used to distinguish experimentally between Bohmian mechanics and standard quantum mechanics. Not surprisingly, an experiment [3] has confirmed standard quantum mechanics.

Struyve *et al.* [4] criticized the conclusion of Golshani and Akhavan. They stressed that, owing to the quantum equilibrium hypothesis [5], it is not possible to achieve the initial condition (2) in a laboratory. Instead, initial positions are distributed according to the quantum mechanical rule

$$\rho(x_1, y_1, x_2, y_2, 0) = \psi^*(x_1, y_1, x_2, y_2, 0)\psi(x_1, y_1, x_2, y_2, 0). \quad (3)$$

The initial distribution (3) implies that particle positions are distributed according to the quantum mechanical rule  $\rho = \psi^*\psi$  at any time  $t$ . In this way, they have shown that there is no incompatibility between Bohmian mechanics and standard quantum mechanics, as far as the situation in [1] is studied.

Recently, Akhavan and Golshani replied to the criticism of Struyve *et al.* by arguing [6] that the initial condition (2) is achieved in a situation in which the initial wavefunction  $\psi(x_1, y_1, x_2, y_2, 0)$  is proportional to

$$\psi(y_1, y_2) = \delta(y_1 - y_2), \quad (4)$$

which, in a similar way, leads to (1) and thus to incompatibility between Bohmian mechanics and standard quantum mechanics. In this short paper we show that this result is an artefact of the fact that the wavefunction (4) is not a regular wavefunction. When the  $\delta$ -function in (4) is regularized in an appropriate way, no incompatibility between Bohmian mechanics and standard quantum mechanics sustains.

For simplicity, we suppress the  $x$ -dependence of the wavefunction. Since the wavefunction  $\psi(y_1, y_2, t)$  satisfies the Schrödinger equation, the density  $\rho(y_1, y_2, t) = \psi^*(y_1, y_2, t)\psi(y_1, y_2, t)$  satisfies the continuity equation

$$\frac{\partial \rho}{\partial t} + \frac{\partial(\rho v_{y_1})}{\partial y_1} + \frac{\partial(\rho v_{y_2})}{\partial y_2} = 0, \quad (5)$$

where  $v_{y_k}(y_1, y_2, t)$  (with  $k = 1, 2$ ) are the Bohmian velocities determined by the wavefunction  $\psi(y_1, y_2, t)$ . The fact that the assumption that particles are initially distributed according to  $\rho(y_1, y_2, 0)$  leads to the result that they are distributed according to  $\rho(y_1, y_2, t)$  at any time  $t$  is a direct consequence of Eq. (5). On the other hand, the result of Akhavan and Golshani [6] corresponds to a situation in which the initial distribution of particles (2) is given by  $\rho(y_1, y_2, 0)$ , but at later times the distribution of particles is not given by  $\rho(y_1, y_2, t)$ . Therefore, their result must be a consequence of a failure in Eq. (5).

It is clear that the failure is due to the use of the wavefunction (4) proportional to a  $\delta$ -function. In fact, the  $\delta$ -function is not a function at all, but rather a functional, meaningful only when integrated over the argument of the functional. Moreover, the square of the  $\delta$ -function, which appears in  $\rho$ , is not well defined even as a functional. Therefore, Eq. (5) does not make sense for the wavefunction proportional to (4). In order to make sense of it, one has to regularize the irregular “function”  $\delta(y)$  (where  $y \equiv y_1 - y_2$ ). The most natural regularization is to use a Gaussian with a small but finite width  $\Delta y$  and then, at the end of calculation, to consider the limit  $\Delta y \rightarrow 0$ . With finite  $\Delta y$ , the initial positions of the particles do not satisfy (2). Instead, they are distributed according to the Gaussian with the width  $\Delta y$ . It is clear that for *any* finite  $\Delta y$ , the distribution of particles at later times will be given by the quantum mechanical distribution  $\rho = \psi^*\psi$ . In that sense, the quantum mechanical distribution at later times is achieved even in the limit  $\Delta y \rightarrow 0$ . Therefore, there is no incompatibility between Bohmian mechanics and standard quantum mechanics.

The source of the wrong conclusion of Akhavan and Golshani in [6] can also be understood in the following qualitative way. Assume that the width of the initial

wavefunction is  $\Delta y_i$ , while that of the final wavefunction is  $\Delta y_f$ . Let their ratio be  $R = \Delta y_f / \Delta y_i$ . If the width of the initial distribution of particles is  $\Delta y_i$ , then the width of the final distribution of particles is equal to the product  $\Delta y_i \cdot R$ . The reasoning in [6] can be reduced to the following reasoning: one studies a case in which  $\Delta y_i = 0$ , so one concludes that the final distribution has the width  $\Delta y_i \cdot R = 0 \cdot R = 0$ . However, in this reasoning, one forgets that  $R = \infty$ . The correct reasoning is  $\Delta y_i \cdot R = 0 \cdot \infty$ , which is indeterminate. The regularization of the indeterminate expression leads to the result that the product is finite and equal to  $\Delta y_f$ , in agreement with standard quantum mechanics.

Finally, note that a wavefunction proportional to a  $\delta$ -function is an idealization that never realizes in nature. Therefore, even if the formal arguments in [6] were correct (which, as argued above, were not), they would not correspond to an incompatibility of Bohmian mechanics with standard quantum mechanics in realistic situations.

This work was supported by the Ministry of Science and Technology of the Republic of Croatia under Contract No. 0098002.

## References

- [1] Golshani M and Akhavan O 2001 *J. Phys. A* **34** 5259
- [2] Bohm D 1952 *Phys. Rev.* **85** 166
- [3] Brida G, Cagliero E, Falzetta G, Genovese M, Gramegna M and Novero C 2002 (*Preprint* quant-ph/0206196)
- [4] Struyve W, De Baere W, De Neve J and De Weirtdt S 2003 *J. Phys. A* **36** 1525
- [5] Dürr D, Goldstein S and Zanghí N 1992 *J. Stat. Phys.* **67** 843
- [6] Akhavan O and Golshani M 2003 (*Preprint* quant-ph/0305020)